

International Congress on **MIDWIFERY AND MATERNAL HEALTH**

April 11, 2022 | Webinar

Labor induction with randomized comparison of cervical, oral and intravaginal misoprostol**Somayeh Fallah and Masoumeh Dadashaliha**

Qazvin University of Medical Science, Iran

Background: This study attempted to evaluate the safety and effectiveness of 50µgm intracervical misoprostol in comparison with intravaginal and sublingual for the induction of labor at term pregnant women.

Methods: This study is designed as a parallel clinical trial study. Three hundred and fifteen term pregnancies requiring induction of labor were treated with the maximum used misoprostol intracervical, sublingual, and vaginal doses. Participants were randomly allocated into three groups of 105. The dose was repeated every 4 hours until adequate uterine contraction and Bishop Score were achieved. The duration of induction to births, time to the active phase, the rate of births, and the need for caesarean section were compared in three groups. Additionally, labor course and side effects were recorded and analyzed. Data were analyzed using SPSS software. A significance level of $p < 0.05$ was considered for statistical analyses.

Findings: Labor was successfully induced in all cases most (63%) of which required a single dose of misoprostol. Ninety-three (93.0%, $p < 0.05$) cervical participants proceeded to vaginal births. This figure was also the same in the vaginal and sublingual group of 83 cases (83.0%). The other 41 cases received caesarean section with more indications of failure to progress and meconium-stained liquor. The results indicated that 278 (92.7%) births were achieved in less than 10 hours. Time from start of medication to the active phase of labor and childbirth was 3.01 ± 0.86 and 6.1 ± 1.3 hours in the Cervical group, 4.2 ± 0.66 and 8.4 ± 0.92 hours in the sublingual group, and 5.06 ± 1.1 and 9.2 ± 1.5 hours in the vaginal group respectively ($p < 0.001$). The Caesarean rate was lower in the cervical group than in the two other groups ($p = 0.05$). No significant differences were observed between the study groups in terms of Apgar score and meconium-stained amniotic fluid. Furthermore, no maternal and neonatal complications were observed.

Conclusion: In addition to the sublingual and intravaginal routes of administration, intracervical misoprostol at a single dose of 50µgm appears to be an effective method for induction of labor in women with an unfavorable cervix. Like all medical interventions, a discussion of the risks, benefits, and alternatives to induction of labor with this medication in each woman should be undertaken before treatment.

Biography

Somayeh Fallah is a Clinical Midwifery Educator and Registered Midwife, she is working more than 10 years in hospital as a staff, instructor as a Faculty member of nursing and midwifery school of Qazvin university of Medical Science in Iran. Moreover, she is a member of Children Growth Research Center, Research Institute for Prevention of Non-Communicable Diseases affiliated to the same university. She is a good researcher, so that she has published many articles that have been presented in many international congresses as a lecturer or poster presentation.